

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07C 311/20, 311/13, 311/28, C07D</b> <b>207/16, 213/40, 213/42, 217/26, 333/38,</b> <b>A61K 31/18</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/37607</b> <b>(43) International Publication Date:</b> 29 July 1999 (29.07.99)
<b>(21) International Application Number:</b> PCT/US99/01663 <b>(22) International Filing Date:</b> 27 January 1999 (27.01.99) <b>(30) Priority Data:</b> 60/072,719 27 January 1998 (27.01.98) US <b>(71) Applicant:</b> ICAGEN, INC. [US/US]; Suite 460, 4222 Emperor Boulevard, Research Triangle Park, NC 27703 (US). <b>(72) Inventors:</b> GROSS, Michael, F.; 1726 Vista Street, Durham, NC 27701 (US). CASTLE, Neil, A.; 116 Canyon Run, Cary, NC 27513 (US). <b>(74) Agents:</b> SKERPON, Joseph, M. et al.; Banner & Witcoff, Ltd., 11th floor, 1001 G Street, N.W., Washington, DC 20001-4597 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (A1, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>With amended claims.</i>

**(54) Title:** POTASSIUM CHANNEL INHIBITORS**(57) Abstract**

Compounds of general formula (I) wherein t is 1 or 2; A and B are each H or taken together form a bond between the substituted carbons; R<sup>1</sup> is H, alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; Y<sup>2</sup> is O, (CH<sub>2</sub>)<sub>q</sub>, HC=CH or NH, w is 0, 1 or 2, q is 0, 1 or 2; X<sup>2</sup> is C=O, C=S or SO<sub>2</sub>; R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroalkyl, an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); Z is in particular H or OH; R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroalkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O- and R<sup>b</sup>R<sup>c</sup>-N-; Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH or ethynyl, p is 0, 1, 2 or 3, o is 0, 1 or 2; X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>, n is 0, 1 or 2; R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroalkyl, an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); or pharmaceutically acceptable salts or prodrugs thereof are useful as potassium channel inhibitors and especially useful for the treatment of cardiac arrhythmias and cell proliferative disorders.



(I)

not a 5-member ring

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



## POTASSIUM CHANNEL INHIBITORS

**BACKGROUND OF THE INVENTION**1. Field of the Invention

The present invention is broadly directed to a class of compounds useful as potassium channel inhibitors.

2. Description of Related Art

Potassium channels are expressed in eukaryotic and procaryotic cells, and are elements in the control of electrical and nonelectrical cellular functions. Subclasses of these channels have been named based on amino acid sequence and functional properties, and include for example voltage gated potassium channels (e.g., Kv1, Kv2, Kv3, Kv4). Subtypes within these subclasses have been characterized as to their putative function, pharmacology and distribution in cells and tissues (Chandy and Gutman, "Voltage-gated potassium channel genes" in Handbook of Receptors and Channels- Ligand and Voltage-gated Ion Channels, ed. R. A. North, 1995; Doupnik et al., *Curr. Opin. Neurobiol.* 5:268, 1995).

Inhibitors of potassium channels lead to a decrease in potassium ion movement across cell membranes. Consequently, such inhibitors induce prolongation of the electrical action potential or membrane potential depolarization in cells containing the inhibited or blocked potassium channels. Prolonging of the electrical action potential is a preferred mechanism for treating certain diseases, e.g., cardiac arrhythmias (Colatsky et al., *Circulation* 82:2235, 1990). Membrane potential depolarization is a preferred mechanism for the treating of certain other diseases, such as those involving the immune system (Kaczorowski and Koo, *Perspectives in Drug Discovery and Design*, 2:233, 1994).

Potassium channels which exhibit functional, pharmacological and tissue distribution characteristics have been cloned. These cloned potassium channels are useful targets in assays for identifying candidate compounds for the treatment of

various disease states. For example, the delayed rectifier voltage-gated potassium channel termed  $I_{Kur}$  or  $I_{Kus}$  which has been reported to contain the Kv1.5  $\alpha$ -subunit gene product is generally believed to be important in the repolarization of the human atrial action potential and thus is a candidate potassium channel target for the treatment of cardiac arrhythmias especially those occurring in the atria (Wang et al., *Circ. Res.* 73:1061, 1993; Fedida et al., *Circ. Res.* 73:210, 1993; Wang et al., *J. Pharmacol. Exp. Ther.* 272:184, 1995; Amos et al., *J. Physiol.*, 491:31, 1996).

The present invention is directed to compounds which are useful as inhibitors of potassium channel function.

It is an object of the present invention, therefore, to provide compounds which are useful for the treatment of diseases in mammals, including humans, and especially for the management of diseases which can be treated by inhibiting cell membrane potassium channels.


Another object of the invention is to provide a method of treating diseases in mammals, including humans, which respond to the inhibition of potassium channel function, which method comprises administering to a mammal in need thereof a compound of the invention.

#### **DETAILED DESCRIPTION OF THE INVENTION**

This invention describes compounds and their utility as inhibitors of potassium channel function. The invention is particularly directed to compounds that inhibit potassium channels which could serve as targets for the treatment of cardiac arrhythmias (i.e.,  $I_{Kur}$ , Kv1.5) especially those occurring in the atria (e.g., atrial flutter and atrial fibrillation) (Wang et al., *Circ. Res.* 73:1061, 1993; Fedida et al., *Circ. Res.* 73:210, 1993; Wang et al., *J. Pharmacol. Exp. Ther.* 272:184, 1995). The present invention also provides a method for treating diseases which respond to the inhibition of potassium channel function. These include, but are not limited to cardiac arrhythmias, cell proliferative disorders including cancer, disorders of the auditory system, central nervous system mediated motor dysfunction and disorders of pulmonary, vascular and visceral smooth muscle contractility.

The invention is particularly based on our discovery that the compounds of the following formula (I) are inhibitors of potassium channel function and are thus useful for inhibiting potassium transport across cellular membranes and for treating cardiac arrhythmias. In particular, these compounds have demonstrated activity against human potassium channels.

Thus, this aspect of the present invention concerns such methods and such compounds having potassium channel inhibitory activity of the formula (I) and pharmaceutically acceptable salts, esters, amides, complexes, chelates, hydrates, stereoisomers, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs thereof.



A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted

heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;


Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^4-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$ .

In another aspect, the present invention concerns such methods and such compounds having potassium channel inhibitory activity of the formula (II) and pharmaceutically acceptable salts, esters, amides, complexes, chelates, hydrates, stereoisomers, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs thereof:



wherein  $t$  is 1, or 2;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ ,  $w$  is 0, 1, or 2 and  $q$  is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$  then  $X^2$  is not  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl;

Z is H, OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;


Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2, or 3; and o is 0, 1, or 2

X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl; and


with the provisos (i) that if Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, p is 0 and X<sup>1</sup> is not (CH<sub>2</sub>)<sub>n</sub>, then R<sup>2</sup> is not H, and (ii) that if R<sup>2</sup> is R<sup>a</sup>-O- and Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub> with p=0,, then X<sup>1</sup> is not SO<sub>2</sub>.

A preferred subgroup of compounds for practicing such methods includes compounds represented by formula (III) and pharmaceutically acceptable salts, esters, amides, complexes, chelates, hydrates, stereoisomers, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs thereof.



wherein t, Y<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as recited above in connection with formula (I), Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, HC=CH, or ethynyl and q is 0, 1, or 2, R<sup>1</sup> is selected from the group of an optionally substituted aryl and an optionally substituted heteroaryl; X<sup>1</sup> is C=O, C=S, or (CH<sub>2</sub>)<sub>n</sub>; wherein n is 0, 1, or 2; and Z is H or OR<sup>14</sup>, where R<sup>14</sup> is H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; where each R<sup>9</sup> is independently selected from H or alkyl; and L is a counter ion.


Another preferred subgroup of compounds for practicing such methods includes compounds represented by formula (IV) and pharmaceutically acceptable salts, esters, amides, complexes, chelates, hydrates, stereoisomers, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs thereof.



wherein t, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as recited above in connection with formula (I), q is 0, 1, or 2, R<sup>1</sup> is H or an optionally substituted aryl selected from the group of phenyl and naphthyl, with the proviso that when q=0, then R<sup>1</sup> cannot be H; X<sup>1</sup> is C=O, or (CH<sub>2</sub>)<sub>n</sub>; Z is H or OH; wherein n is 0, 1, or 2; and Y<sup>1</sup> is CH=CH, ethynyl, or (CH<sub>2</sub>)<sub>p</sub>; where p is 0, 1, 2 or 3.

A particularly preferred subgroup of compounds for practicing such methods

includes compounds represented by formula (V) and pharmaceutically acceptable salts, esters, amides, complexes, chelates, hydrates, stereoisomers, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs thereof:




wherein  $R^2$ ,  $R^3$  and  $R^4$  are as recited above in connection with formula (I) ( $R^3$  preferably is H), where  $R^1$  is an optionally substituted aryl selected from the group of phenyl and naphthyl; Z is H, or OH;  $X^1$  is C=O, or  $(CH_2)_n$ ; wherein n is 0, 1, or 2; and  $Y^1$  is CH=CH, ethynyl or  $(CH_2)_p$ , where p is 0, 1, 2 or 3.

In the above formulae,  $R^1$  and  $R^2$  are preferably moieties that are non-ionized at a physiological pH. In preferred aspects of the present invention,  $R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl and where  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl and  $R^3$  and  $R^4$  are independently selected from H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl in the above formulae (I), (II), (III), (IV) and (V). Compounds according to the present invention are particularly directed to those compounds of formulae (I), (II), (III), (IV) and (V) subject to the proviso that when  $R^1$  is an optionally substituted aryl, then said optionally substituted aryl is not a dialkoxyphenyl, and especially is not a 3,4-dialkoxyphenyl.

Further preferred compounds are those having the previously identified



formulae (I) (where A and B are hydrogen), (II), (III), (IV), or (V); but having the stereochemical configuration of substituents attached to the saturated ring of the core structure in accordance with the following representative formula (VI):



Still other preferred compounds of the present invention are those of formulae (I), (II), (IV) and (VI) having the ring substituents in the orientation of previous formulae (III) and (V).

The term "alkyl" as used alone or in combination herein refers to a straight or branched chain saturated hydrocarbon group containing from one to ten carbon atoms. Preferably, the alkyl group is a "C<sub>1-6</sub> alkyl" or "lower alkyl" which refer to such groups containing from one to six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl and the like.

The related term "alkylene," as used alone or in combination herein, refers to a straight or branched chain saturated divalent hydrocarbon group containing from one to ten carbon atoms. Preferably, the alkylene group is a "C<sub>1-6</sub> alkylene" or "lower alkylene" which refer to such groups containing from one to six carbon atoms, such as methylene, ethylene, n-propylene, isopropylene, n-butylene, isobutylene, sec-butylene, tert-butylene and the like.

The term "alkoxy" as used alone or in combination herein refers to a straight or branched chain alkyl group covalently bonded to the parent molecule through an -O- linkage containing from one to ten carbon atoms and the terms "C<sub>1-6</sub> alkoxy" and "lower alkoxy" refer to such groups containing from one to six carbon atoms, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy and the like.

The term "alkoxyalkyl" refers to an alkyl group substituted with an alkoxy

group.

The term "haloalkyl" is a substituted alkyl, preferably a substituted lower alkyl, substituted with one or more halogen atoms, and preferably is a C<sub>1</sub> to C<sub>4</sub> alkyl substituted with one to three halogen atoms. One example of a haloalkyl is trifluoromethyl.

The term "alkanoyl" as used alone or in combination herein refers to an acyl radical derived from an alkanecarboxylic acid, particularly a lower alkanecarboxylic acid, and includes such examples as acetyl, propionyl, butyryl, valeryl, and 4-methylvaleryl.

The term "aminocarbonyl" means an amino-substituted carbonyl (carbamoyl or carboxamide) wherein the amino group can be a primary, secondary (mono-substituted amino) or tertiary amino (di-substituted amino) group preferably having as a substituent(s) a lower alkyl.

The term "carbocycloalkyl" refers to stable, saturated or partially unsaturated monocyclic, bridged monocyclic, bicyclic, and spiro ring hydrocarbyls of 3 to 15 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclohexyl, bicyclooctyl, bicyclononyl, spirononyl and spirodecyl. The term "optionally substituted" as it refers to "carbocycloalkyl" herein indicates that the carbocycloalkyl group may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups.

The term "heterocyclyl" as used herein refers to a stable, saturated, or

partially unsaturated, monocyclic, bridged monocyclic, bicyclic, and spiro ring system containing carbon atoms and other atoms selected from nitrogen, sulfur and/or oxygen. Preferably, a heterocyclyl is a 5 or 6-membered monocyclic ring or an 8-11 membered bicyclic ring which consists of carbon atoms and contains one, two, or three heteroatoms selected from nitrogen, oxygen and/or sulfur. The term "optionally substituted" as it refers to "heterocyclyl" herein indicates that the heterocyclyl group may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Examples of such heterocyclyl groups are isoxazolyl, imidazolyl, thiazolyl, imidazolidinyl, pyrrolyl, pyrrolinyl, pyranyl, pyrazinyl, piperidyl, morpholinyl and triazolyl. The heterocyclyl group may be attached to the parent structure through a carbon atom or through any heteroatom of the heterocyclyl that results in a stable structure.

The term "heteroaryl" as used herein refers to a stable, aromatic monocyclic or bicyclic ring system containing carbon atoms and other atoms selected from nitrogen, sulfur and/or oxygen. Preferably, a heteroaryl is a 5 or 6-membered monocyclic ring (optionally benzofused) or an 8-11 membered bicyclic ring which consists of carbon atoms and contains one, two, or three heteroatoms selected from nitrogen, oxygen and/or sulfur. The term "optionally substituted" as it refers to "heteroaryl" herein indicates that the heteroaryl group may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro,

monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino, cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Examples of such heteroaryl groups are isoxazolyl, imidazolyl, thiazolyl, isothiazolyl, pyridyl, furyl, pyrimidinyl, pyrazolyl, pyridazinyl, furazanyl and thienyl. The heteroaryl group may be attached to the parent structure through a carbon atom or through any heteroatom of the heteroaryl that results in a stable structure.

The term "heteroaralkyl" as used herein refers to a lower alkyl as defined above in which one hydrogen atom is replaced by a heteroaryl radical as defined above. The term "optionally substituted" as it refers to "heteroaralkyl" herein indicates that the heteroaryl group may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino, cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Examples of such heteroaralkyl groups are 2-pyridylmethyl, 3-pyridylmethyl, 4-pyridylmethyl, 3-pyridylethyl and 4-pyrimidinylmethyl.

The specific chemical nature of the optionally substituted heterocyclyl and heteroaryl groups for the terminal moieties R<sup>1</sup> and R<sup>2</sup> in the prior identified potassium channel inhibitor compounds is not narrowly critical and, as noted above, a wide

variety of substituent groups are contemplated. Preferably, the substituents for the heterocyclyl and heteroaryl groups are selected such that the total number of carbon and hetero atoms comprising the substituted heterocyclyls and heteroaryls is no more than about 25.

5           The terms "halo" and "halogen" as used herein to identify substituent moieties, represent fluorine, chlorine, bromine or iodine, preferably chlorine or fluorine.

10           The term "aryl" when used alone or in combination refers to an unsubstituted or optionally substituted monocyclic or bicyclic aromatic hydrocarbon ring system. Preferred are optionally substituted phenyl or naphthyl groups. The aryl group may optionally be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Preferably, the aryl group is phenyl optionally substituted with up to four and usually with one or two groups, preferably selected from C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, as well as cyano, trifluoromethyl and halo.

20           The term "aralkyl" alone or in combination refers to a lower alkyl radical as defined above in which one hydrogen atom is replaced by an aryl radical as defined above, and includes benzyl, and 2-phenylethyl. The aralkyl group may optionally be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), aralkyl, alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl),

alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups.

The term "alkoxycarbonyl" alone or in combination means a radical of the formula -C(O)-alkoxy, in which alkoxy is as defined above.

The term "alkylcarbonyloxy" alone or in combination means a radical of the formula -O-C(O)-alkyl, in which alkyl is as defined above.

The term "alkenyl" means a two to seven carbon, straight or branched hydrocarbon containing one or more double bonds, preferably one or two double bonds. Examples of alkenyl include ethenylene, propenylene, 1, 3- butadienyl, and 1, 3, 5-hexatrienyl.

The term "substituted amino" refers to a group of the formula NZ'Z" wherein Z' is H, alkyl, carbocycloalkyl, aryl, heteroaryl, heterocyclyl, heteroaralkyl, or heterocyclyl(alkylene) and Z" is H, alkyl, carbocycloalkyl, or aryl further substituted with a carboxylic acid or carboxylic ester, provided that when Z' is H, then Z" is other than H, or Z' and Z" taken together with the nitrogen atom to which they are attached are 1-pyrrolidinyl, 1-piperidinyl, 1-azepinyl, 4-morpholinyl, 4-thiamorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, each optionally substituted with alkyl, alkoxy, alkylthio, halo, aryl or hydroxy.

The term "treating" as used herein, describes the management and care of a patient afflicted with a condition, disease or disorder for which the administration of a compound of the present invention alters the action or activity of a potassium channel to prevent the onset of symptoms or complications associated with the condition, disease or disorder, to alleviate the symptoms or complications caused by the condition, disease or disorder, or to eliminate the condition, disease or disorder altogether.

It is recognized that there may be one or two chiral centers in the compounds falling within the scope of the present invention and thus such compounds will exist as various stereoisomeric forms. Applicants intend to include all the various stereoisomers within the scope of the invention, referred to herein as the  
5 "pharmaceutically acceptable stereoisomers." Thus, this invention is intended to include the cis and trans isomers and the corresponding enantiomers of the compounds of formula I-IV. Though the compounds may be prepared as racemates and can conveniently be used as such, individual enantiomers also can be isolated or preferentially synthesized by known techniques if desired. Such racemates and  
10 individual enantiomers and mixtures thereof are intended to be included within the scope of the present invention.

The present invention also encompasses the pharmaceutically acceptable esters, amides, complexes, chelates, hydrates, crystalline or amorphous forms, metabolites, metabolic precursors or prodrugs of the compounds of formulae (I), (II),  
15 (III) and (IV). Pharmaceutically esters and amides can be prepared by reacting, respectively, a hydroxy or amino functional group with a pharmaceutically acceptable organic acid, such as identified below. A prodrug is a drug which has been chemically modified and may be biologically inactive at its site of action, but which is degraded or modified by one or more enzymatic or other *in vivo* processes to the parent  
20 bioactive form. Generally, a prodrug has a different pharmacokinetic profile than the parent drug such that, for example, it is more easily absorbed across the mucosal epithelium, it has better salt formation or solubility and/or it has better systemic stability (e.g., an increased plasma half-life).

Those skilled in the art recognize that chemical modifications of a parent drug to yield a prodrug include: (1) terminal ester or amide derivatives which are  
25 susceptible to being cleaved by esterases or lipases; (2) terminal peptides which may be recognized by specific or nonspecific proteases; or (3) a derivative that causes the prodrug to accumulate at a site of action through membrane selection, and combinations of the above techniques. Conventional procedures for the selection and

preparation of prodrug derivatives are described in H. Bundgaard, *Design of Prodrugs*, (1985). Those skilled in the art are well-versed in the preparation of prodrugs and are well-aware of its meaning.

5 The compounds of the present invention can be used in their neat form or in the form of pharmaceutically-acceptable salts derived from inorganic or organic acids. Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts of compounds of the present invention include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid. These salts thus include, but are not  
10 limited to, the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate,  
15 methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate.

Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates, like dimethyl, diethyl, dibutyl and diamyl  
20 sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, omides and iodides, aralkyl halides like benzyl and phenethyl bromides and others. Water or oil soluble or dispersible products are thereby generally obtained.

25 The pharmaceutically acceptable salts of the compounds of the present invention also can exist as various solvates, such as with water, methanol, ethanol, dimethylformamide, ethyl acetate and the like. Mixtures of such solvates also can be prepared. Such solvates are within the scope of the present invention.

The pharmacological profile of the potassium channel inhibitory activity of the compounds of the present invention can be readily assessed by those skilled in the art



5 using routine experimentation, such as the procedures and techniques illustrated in the examples which follow. Assays for assessing the activity of particular compounds may employ cells stably transfected to express a specific potassium channel, as well as native mammalian cells. In particular, cells stably transfected to express a specific potassium channel, which have been treated with a voltage dependent fluorescent dye, such as *bis*-(1,3-dibutylbarbituric acid)trimethine oxonol, can be used to gauge the inhibitory activity of potassium channel inhibitor compounds, possibly in comparison to known inhibitors. Alternatively, such cells can be primed with a detectible species, such as  $^{86}\text{Rb}$ , and then challenged with a particular compound, under conditions otherwise suitable for activating the potassium channel, to assess the potassium inhibitory activity of the compound. The potassium channel inhibitory activity of a compound also can be determined using isolated mammalian cells and the whole cell configuration of the known patch clamp technique (Hamill et al., *Pflugers Archiv* 391:85, 1981). These and other known techniques can be readily employed by those skilled in the art to assess the activity level of the potassium channel inhibitor compounds of the present invention.

10 The compounds of the present invention may be administered by a variety of routes including orally, parenterally, sublingually, intranasally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intracardiac injection, or infusion techniques. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices.

25 Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,2-propanediol. Among the

acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed as mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and phosphatidyl cholines (lecithins), both

natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33, et seq.

5 To select preferred compounds from less preferred compounds, one uses by example the *in vitro* assays detailed under the sub-heading **BioAssays** hereafter. Typically, a preferred compound will produce half maximal blocking activity at a concentration ranging from about 10nM to about 1 $\mu$ M in the *in vitro* assays described. One of ordinary skill will recognize that the final and optimum dose and regimen will be determined empirically for any given drug.

10 Total daily dose administered to a host in single or divided doses may be an amount, for example, from 0.001 to 100 mg of active ingredient per kg body weight on a daily basis and more usually 0.01 to 10 mg/kg/day. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose. It is anticipated that a therapeutically effective serum concentration of active ingredient  
15 will be 10 nM to 10 $\mu$ M (5ng/ml to 5 $\mu$ g/ml).

The amount of active ingredient that may be combined with carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

20 It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, and diet of the patient, the time of administration, the route of administration, the rate of excretion, whether a drug combination is used, and the severity of the particular disease.

25 The present invention is explained in greater detail in the Examples which follow. These examples are intended as illustrative of the invention, and are not to be taken as limiting thereof. Unless otherwise indicated, all references to parts and percentages are based on weight and all temperatures are expressed in degrees Celsius. The scope of the invention is not construed as merely consisting of the following examples.

## EXAMPLES

Unless otherwise specified, all solvents and reagents were purchased from commercial suppliers and used without further purification. Analytical thin layer chromatography (TLC) was performed on Whatman Inc. 60 silica gel plates (0.25 mm thickness). Compounds were visualized under UV lamp or by developing with  $\text{KMnO}_4/\text{KOH}$ , ninhydrin, or Hanessian's solution. Flash chromatography was done using silica gel from Selectro Scientific (particle size 32-63).  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded at 300 MHz and 75.5 MHz, respectively.


### Compound Preparation

Tetrahydronaphthalene (tetralin) and benzocycloheptane, compounds of the previous formulae (I), (II), (III) and (IV) useful as potassium channel inhibitors in accordance with the present invention can be prepared in accordance with several sequential steps as illustrated with reference to the tetralin species in the preparation which follow.

#### Preparation 1

##### Synthesis of 7-nitro-1,2,3,4-tetrahydro-2-naphthalenol

This preparation demonstrates the reduction of a nitrotetralone to give the corresponding alcohol.




A suspension of 7-nitro-1-tetralone (10.14 g, 0.053 mol) in  $\text{MeOH}$  (600 ml) was cooled to  $0^\circ\text{C}$  and treated with  $\text{NaBH}_4$  (4.25 g, 0.11 mol, 2.1 equiv.). A nitrotetralone can be obtained by nitration of a 1-tetralone, the desired product being separated from minor component byproducts. The reaction mixture became homogeneous almost immediately. After stirring at  $0^\circ\text{C}$  for 30 min, 2N  $\text{HCl}$  (100 ml) was added and stirring was continued for an additional 30 min. The reaction mixture was concentrated under reduced pressure (approx. 150 ml) and diluted with  $\text{CH}_2\text{Cl}_2$  (200 ml) and  $\text{H}_2\text{O}$  (100 ml). The aqueous layer was separated and extracted with additional  $\text{CH}_2\text{Cl}_2$  (2 x 100 ml). The combined

organic layers were washed with brine (100 ml), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated under reduced pressure to 7-nitro-1,2,3,4-tetrahydro-2-naphthalenol as a white solid (10.13 g, 99%) which was used in the next step without further purification.  $R_f$  (silica gel): 0.50 (40% hexane: 40%  $\text{CH}_2\text{Cl}_2$ : 20% EtOAc);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) 8.29 (d,  $J = 2.1$  Hz, 1H), 7.97 (dd,  $J = 2.1$  and 8.1 Hz, 1H), 7.21 (d,  $J = 8.1$  Hz, 1H), 4.80–4.77 (m, 1H), 2.94–2.73 (m, 2H), 2.47 (d,  $J = 6.0$  Hz, 1H), 2.12–1.93 (m, 2H), 1.90–1.74 (m, 2H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ) 146.5, 145.1, 140.6, 129.9, 123.6, 122.2, 67.8, 31.9, 29.3, 18.6.

### Preparation 2

#### Synthesis of 7-nitro-3,4-dihydronaphthalene

This preparation describes subjecting the alcohol product of Preparation 1 to an acid catalyzed dehydration to give the corresponding tetralene.




7-nitro-1,2,3,4-tetrahydro-2-naphthalenol (10.13 g, 0.053 mol) (from Preparation 1) was heated in the presence of TsOH- $\text{H}_2\text{O}$  (1.72 g, 0.009 mol, 0.2 equiv.) in toluene (150 ml) for 2 h at 100 °C. The solvent was removed under reduced pressure and the residue was treated with EtOAc (150 ml) and saturated aqueous  $\text{NaHCO}_3$  (150 ml). The aqueous layer was separated and extracted with additional EtOAc (2 x 100 ml). The combined organic layers were washed with saturated aqueous NaCl (200 ml), dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated under reduced pressure to give 7-nitro-3,4-dihydronaphthalene as a brown oil (9.18 g, 100%) which was used in the next step without additional purification.  $R_f$  (silica gel): 0.79 (70% hexane: 30% EtOAc);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) 7.95 (dd,  $J = 2.4$  and 8.1 Hz, 1H), 7.83 (d,  $J = 2.4$  Hz, 1H), 7.21 (d,  $J = 8.1$  Hz, 1H), 6.50 (d,  $J = 6.50$  Hz, 1H), 6.18 (dt,  $J = 4.5$  and 9.6 Hz, 2H), 2.88 (t,  $J = 8.4$  Hz, 2H), 2.40–2.34 (m, 2H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ) 147.1, 143.1, 135.3, 131.4, 128.2, 126.5, 121.8, 120.3, 27.4, 22.5.

### Preparation 3

#### Synthesis of 1,2-epoxy-7-nitro-3,4-dihydronaphthalene

In this preparation, the double bond in the tetralene of Preparation 2 is oxidized to give the corresponding epoxide.




A solution of 7-nitro-3,4-dihydronaphthalene (9.18g; 0.052 mol) (from Preparation 2) in CH<sub>2</sub>Cl<sub>2</sub> (600 ml) was cooled to 0 °C and treated with *m*-CPBA, 57-85%, (13.86 g, approx. 0.056 mol, approx. 1.1 equiv). The reaction mixture was allowed to stir for 48 h, slowly warming to room temperature. The mixture was treated with aqueous NaHCO<sub>3</sub> (300 ml) and the organic layer was separated. The organic layer was extracted with additional aqueous NaHCO<sub>3</sub>, washed with aqueous NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated under reduced pressure to give 1,2-epoxy-7-nitro-3,4-dihydronaphthalene (9.94, 100%) as a white solid which was used in the next step without further purification. . R<sub>f</sub> (silica gel): 0.56 (70% hexane: 30% EtOAc); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) 8.24 (s, 1H), 8.08 (dd, *J* = 1.8 and 8.1 Hz, 1H), 7.23 (d, *J* = 8.1 Hz, 1H), 3.92 (d, *J* = 4.2 Hz, 1H), 3.77 (s, 1H), 2.87-2.62 (m, 2H), 2.47 (dd, *J* = 6.6 and 14.4 Hz, 1H), 1.78 (dt, *J* = 5.7 and 14.1 Hz, 1H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) 146.5, 144.7, 134.5, 129.4, 124.4, 123.4, 54.7, 51.8, 24.5, 21.0.

### Preparation 4

#### Synthesis of *trans*-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol

In this preparation, the epoxide is reacted with ammonium hydroxide to give the corresponding amino alcohol.




A solution of 1,2-epoxy-7-nitro-3,4-dihydronaphthalene (10.84 g; 0.057 mol) (from Preparation 3) in THF (50 ml) and EtOH (50 ml) was heated to 40°C and NH<sub>4</sub>OH (60 ml) was added dropwise over the course of 1 h. After the addition was complete, the

temperature was increased to 60°C and the reaction was stirred for 24 h. An additional 50 ml of  $\text{NH}_4\text{OH}$  was added and the reaction was stirred for another 24 h. The solvent was removed under reduced pressure to give a brown powder (10.73 g) that was dried under high vacuum at 50°C for 48 h. The *trans*-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol was used in the next two preparations without further purification.

#### Preparation 5

The general synthesis of secondary amines as illustrated for the synthesis of give *trans*-*N*-(benzyl)-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol.

In this preparation, the amino alcohol is reacted with an aldehyde to attach an  $\text{R}'$ -moiety to the amino group, where  $\text{R}'$  is equivalent to  $\text{R}^3$  as defined in formula (I).



The amino alcohol is reacted in a suitable solvent with the aldehyde under reductive amination conditions. Suitable solvents in which the reaction can be conducted include glacial acetic acid,  $\text{MeOH}$ , or 1,2-dichloroethane. Suitable reducing agents include sodium triacetoxyborohydride, sodium cyanoborohydride, or sodium borohydride.

A solution of *trans*-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol (0.58 g; 2.8 mmol) (from Preparation 4) in glacial acetic acid was treated with benzaldehyde (0.31 ml; 3.0 mmol; 1.1 equiv.) followed by sodium triacetoxyborohydride (0.82 g; 3.9 mmol; 1.4 equiv.). The reaction mixture was allowed to stir at room temperature for 16 h. The reaction mixture was diluted with  $\text{EtOAc}$  (50 ml) and the pH was adjusted to  $\text{pH} = 9$  by the addition of 1 N  $\text{NaOH}$ . The organic layer was separated, washed with aqueous  $\text{NaCl}$  (50 ml), filtered, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The crude product was purified by flash chromatography on silica gel to give *trans*-*N*-(benzyl)-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol (0.37 g; 44%).  $R_f$  (silica gel)


0.58 (60% EtOAc: 20% hexane: 20% CH<sub>2</sub>Cl<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-acetone) 8.38 (d, *J* = 2.1 Hz, 1H), 7.96 (dd, *J* = 2.1 and 8.7 Hz, 1H), 7.43 (d, *J* = 7.2 Hz, 2H), 7.35-7.30 (m, 3H), 7.25-7.20 (m, 1H), 4.20-4.14 (m, 1H), 3.93 (d, *J* = 13.5 Hz, 1H), 3.81 (s, 1H), 3.80 (d, *J* = 6 Hz, 1H), 3.77 (d, *J* = 13.5 Hz, 1H), 3.07-2.84 (m, 2H), 2.27-2.17 (m, 1H), 1.97-1.96 (m, 1H); <sup>13</sup>C NMR (75 MHz, d<sub>6</sub>-acetone) 146.5, 145.6, 141.3, 139.8, 129.6, 128.3 (two carbons), 128.2 (two carbons), 126.8, 124.2, 121.1, 67.2, 61.9, 49.8, 27.5, 26.3.

### Preparation 6

The general procedure for the synthesis of sulfonamides as illustrated for the synthesis of *trans*-*N*-(4-ethylphenylsulfonyl)-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol.

While the amino alcohol of Preparation 4 or 5 can be optionally protected with conventional protecting group(s) as are commonly employed to block or protect the amino (-NH<sub>2</sub>) and/or the hydroxy (-OH) functionality while reacting other functional groups on the parent compound, this (and the subsequent) preparations shows that it is possible to react the amino alcohol directly without use of any protecting group(s).

In this preparation, the amino alcohol is reacted with a sulfonyl chloride to attach an R'-SO<sub>2</sub>- moiety to the amino group, where R' is equivalent to R<sup>1</sup>-Y<sup>2</sup> as defined in formula (I) and elsewhere. The amino alcohol is reacted in a suitable solvent with the sulfonyl chloride (R'SO<sub>2</sub>Cl) or sulfonyl anhydride in the presence of an acid scavenger. Suitable solvents in which the reaction can be conducted include methylene chloride, DMF and tetrahydrofuran. Suitable acid scavengers include triethylamine, and pyridine.



A solution of *trans*-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol (0.91 g; 4.37 mmol) (from Preparation 4) in THF (20 ml) was cooled to 0 °C and treated with




DMAP (0.010 g; 0.082 mmol, 0.02 equiv), NEt<sub>3</sub> (0.90 ml; 6.46 mmol; 1.5 equiv) and 4-ethylbenzene sulfonyl chloride (1.05 g; 5.13 mmol; 1.2 equiv). After 15 min at 0°C, the reaction was allowed to warm up to room temperature and stirred for an additional 24 h. The solvent was removed under reduced pressure and the residue was treated with EtOAc (150 ml) and a 20% aqueous solution of conc. HCl (50 ml). The organic layer was separated, washed with aqueous NaCl (50 ml), filtered, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated. The crude product was purified by flash chromatography on silica gel to give *trans-N*-(4-ethylphenylsulfonyl)-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol as a tan solid (1.10 g; 67%). R<sub>f</sub> (silica gel): 0.67 (60% EtOAc: 20% hexane: 20% CH<sub>2</sub>Cl<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) 7.93 (d, *J* = 2.1 Hz, 1H), 7.89 (d, *J* = 8.4 Hz, 2H), 7.44 (d, *J* = 8.4 Hz, 2H), 7.40 (d, *J* = 2.1 Hz, 1H), 7.20 (d, *J* = 8.7 Hz, 1H), 5.21 (d, *J* = 8.1 Hz, 1H), 4.28 (t, *J* = 7.8 Hz, 1H), 4.07-4.04 (m, 1H), 3.03 (d, *J* = 2.7 Hz, 1H), 2.98-2.88 (m, 2H), 2.77 (q, *J* = 7.5 Hz, 2H), 2.51-2.16 (m, 1H), 1.29 (t, *J* = 7.5 Hz, 3H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) 150.7, 146.5, 144.7, 136.9, 135.7, 129.7, 129.1 (two carbons), 127.1 (two carbons), 123.8, 122.3, 70.4, 58.4, 28.6, 27.1, 26.3, 14.7.

#### Preparation 7

The general procedure for the reduction of the aromatic nitro functionality as illustrated for the synthesis of *trans-N*-(4-ethylphenylsulfonyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol.

The sulfonylated product of Preparation 6 is reduced in this preparation to give the corresponding aniline.




A solution of *trans-N*-(4-ethylphenylsulfonyl)-1-amino-7-nitro-1,2,3,4-tetrahydro-2-naphthalenol (0.96 g; 2.6 mmol) (from Preparation 6) in THF (15 ml) and MeOH (10 ml) was cooled to 0°C and treated with NaBH<sub>4</sub> (0.46 g; 12.2 mmol; 4.7 equiv.) followed

immediately by  $\text{NiCl}_2$  (0.15 g; 1.2 mmol, 0.5 equiv.). After 15 min at  $0^\circ\text{C}$ , the reaction was allowed to warm up to room temperature and stirred for an additional 1 h. The solvent was removed under reduced pressure to leave a black residue which was treated with EtOAc (100 ml) and aqueous NaCl (100 ml). The aqueous layer was separated and extracted with additional EtOAc (3 x 50 ml). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated under reduced pressure to give *trans*-N1-(4-ethylphenylsulfonyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol as a tan solid (0.79g; 89%) which was used without further purification in the next step.  $R_f$  (silica gel): 0.43 (60% EtOAc: 20% hexane: 20%  $\text{CH}_2\text{Cl}_2$ );  $^1\text{H}$  NMR (300 MHz,  $d_4$ -MeOH) 7.86 (d,  $J = 8.4$  Hz, 2H), 7.45 (d,  $J = 8.4$  Hz, 2H), 6.80 (d,  $J = 8.1$  Hz, 1H), 6.55 (dd,  $J = 2.4$  and 8.4 Hz, 1H), 6.03 (d,  $J = 1.8$  Hz, 1H), 4.10 (d,  $J = 4.8$  Hz, 1H), 3.92-3.88 (m, 1H), 2.81-2.71 (m, 3H), 2.60-2.51 (m, 1H), 2.06-1.96 (m, 1H), 1.81-1.73 (m, 1H), 1.30 (t,  $J = 7.5$  Hz, 3H);  $^{13}\text{C}$  NMR (75 MHz,  $d_6$ -DMSO) 148.8, 146.8, 140.8, 135.5, 129.1, 128.8, 127.1, 124.5, 115.5, 114.5, 68.4, 57.1, 28.4, 25.9, 23.4, 15.5.

#### Preparation 8

##### Synthesis of *trans*-N1-(4-ethylphenylsulfonyl)-N7-(4-trifluoromethoxybenzyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol

In this preparation, the amino group on the aniline product of Preparation 7 is substituted.




A solution of *trans*-N1-(4-ethylphenylsulfonyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol (0.049 g; 0.14 mmol) (from Preparation 6) in anhydrous DMF (2 ml) was treated with  $\text{K}_2\text{CO}_3$  (0.040 g; 0.29 mmol; 2.1 equiv.) and 18-crown-6 (0.060 g; 0.23 mmol; 1.6 equiv.) followed by 4-trifluoromethoxybenzyl bromide (30  $\mu\text{M}$ ; 0.19 mmol;

1.3 equiv.). The reaction mixture was heated to 60°C and allowed stir 24 h. The reaction mixture was diluted with EtOAc (10 ml) and 1N HCl (20 ml). The organic layer was separated, washed with additional 1N HCl (20 ml) and brine (20 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was purified by flash chromatography on silica gel to give *trans*-N1-(4-ethylphenylsulfonyl)-N7-(4-trifluoromethoxybenzyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol (0.035 g; 48%) as a white solid. R<sub>f</sub> (silica gel): 0.54 (30% EtOAc: 40% hexane: 30% CH<sub>2</sub>Cl<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) 7.84 (d, *J* = 7.8 Hz, 1H), 7.77 (d, *J* = 8.1 Hz, 2H), 7.39 (d, *J* = 6.0 Hz, 2H), 7.36 (d, *J* = 6.0 Hz, 2H), 7.28 (d, *J* = 8.4 Hz, 2H), 6.70 (d, *J* = 8.4 Hz, 1H), 6.39 (dd, *J* 2.1 and 8.4 Hz, 1H), 6.03 (d, *J* = 1.8 Hz, 1H), 5.89 (t, *J* = 6.0 Hz, 1H), 4.68 (d, *J* = 3.3 Hz, 1H), 4.03-3.88 (m, 3H), 3.68 (d, *J* = 3.3 Hz, 1H), 2.58 (q, *J* = 7.5 Hz, 2H), 2.64-2.57 (m, 1H), 2.39-2.30 (m, 1H), 1.91-1.81 (m, 1H), 1.60-1.54 (m, 1H), 1.12 (t, *J* = 7.5 Hz, 3H); <sup>13</sup>C NMR (75 MHz, d<sub>6</sub>-DMSO) 148.8, 147.7, 146.9, 140.5, 140.4, 135.3, 129.7, 129.3, 128.8, 127.2, 124.9, 121.4, 113.5, 113.0, 68.8, 57.0, 46.2, 28.3, 25.6, 23.2, 15.3.

### Preparation 9

#### Synthesis of *trans*-N1-(4-*n*-propylphenylsulfonyl)-N7-(styrylcarbamoyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol

In this preparation, an aniline analogous to that of Preparation 7 is acylated, for example using RCOCl where R is equivalent to R<sup>2</sup>-Y<sup>1</sup> and X<sup>1</sup> is C=O as defined in formula (I) and elsewhere to attach a substituent group to the amino group.






A solution of *trans*-N1-(4-*n*-propylphenylsulfonyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol (0.076 g; 0.21 mmol) in anhydrous DMF (2 ml) was cooled to 0°C and treated with NEt<sub>3</sub> (30 μL, 0.22 mmol; 1 equiv) followed by cinnamoyl chloride (0.049


g; 0.29 mmol; 1.4 equiv). After 15 min at 0°C, the reaction was allowed to warm up to room temperature and stirred for an additional 12 h. The reaction mixture was diluted with EtOAc (15 ml) and 1N HCl (20 ml). The organic layer was separated, washed with additional 1N HCl (20 ml) and brine (20 ml), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was purified by flash chromatography on silica gel to give *trans*-N1-(4-*n*-propylphenylsulfonyl)-N7-(styrylcarbamoyl)-1,7-diamino-1,2,3,4-tetrahydro-2-naphthalenol (0.061 g; 59%) as a white solid. R<sub>f</sub> (silica gel): 0.61 (60% EtOAc: 20% hexane: 20% CH<sub>2</sub>Cl<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, *d*<sub>6</sub>-DMSO) 10.06 (s, 1H), 7.96 (d, *J* = 8.1 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 2H), 7.62-7.53 (m, 4H), 7.46-7.39 (m, 5H), 7.04 (d, *J* = 8.4 Hz, 1H), 6.82 (d, *J* = 15.6 Hz, 1H), 4.81 (d, *J* = 3.0 Hz, 1H), 4.16 (d, *J* = 4.1 Hz, 1H), 3.61 (d, *J* = 2.7 Hz, 1H), 2.76-2.66 (m, 1H), 2.61 (t, *J* = 7.5 Hz, 2H), 2.58-2.50 (m, 1H), 2.00-1.85 (m, 1H), 1.60-1.53 (m, 3H), 0.85 (t, *J* = 7.5 Hz, 3H); <sup>13</sup>C NMR (75 MHz, *d*<sub>6</sub>-DMSO) 163.8, 147.2, 140.6, 140.4, 137.5, 135.6, 135.4, 132.5, 130.2, 129.6 (two carbons), 129.4 (two carbons), 129.1, 128.2 (two carbons), 126.9 (two carbons), 123.0, 121.9, 119.4, 67.9, 56.8, 37.4, 25.2, 24.0, 23.5, 13.9.

When using a protecting group in connection with a specific synthesis, the species of protecting group used is not critical so long as the derivatized -NH<sub>2</sub> or -OH group is stable to the condition(s) of subsequent reaction(s) and can be removed at the appropriate point without disrupting the remainder of the molecule. For amino protecting groups see T.W. Greene and P. Wuts, Protective Groups in Organic Synthesis, Chapter 7 (1991). Preferred amino protecting groups are t-butoxycarbonyl (Boc), phthalimide, a cyclic alkyl, and benzyloxycarbonyl. For hydroxy protecting groups see T.W. Greene and P. Wuts, Protective Groups in Organic Synthesis, Chapter 2 (1991). A suitable "hydroxy protecting group" includes one of the ether or ester derivatives of the hydroxy group commonly employed to block or protect the hydroxy group while reactions are carried out on other functional groups on a compound. Hydroxy protecting groups include tert-butyldiphenylsilyloxy (TBDPS), tert-butyldimethylsilyloxy (TBDMS), triphenylmethyl (trityl), mono- or di-methoxytrityl, or an alkyl or aryl ester.


Using the principles and techniques of Preparations 1 through 9 (and methods available from the literature, such as WO 98/04521 and WO 98/36749), and appropriate starting materials, which will be well-understood by those skilled in the art, a variety of other compounds falling within the scope of the present invention can be synthesized. In this regard, compounds listed in the following Tables 1A, 1B and 1C can be synthesized.

Table 1A


			
Entry	R <sup>1</sup>	R <sup>2</sup>	-Y <sup>1</sup> -X <sup>1</sup> -
1	4-ethylphenyl	3-methoxyphenyl	-C(O)-
2	3-ethylphenyl	3-methoxyphenyl	-C(O)-
3	3-n-propylphenyl	3-methoxyphenyl	-C(O)-
4	4-ethylphenyl	4-methoxyphenyl	-C(O)-
5	4-ethylphenyl	4-chlorophenyl	-C(O)-
6	4-ethylphenyl	4-pentoxyphenyl	-C(O)-
7	4-isopropylphenyl	3-methoxyphenyl	-C(O)-
8	4-n-propylphenyl	3-methoxyphenyl	-C(O)-
9	4-ethylphenyl	3-ethoxyphenyl	-C(O)-
10	4-chlorophenyl	3-ethoxyphenyl	-C(O)-
11	4-chlorophenyl	phenyl	-C(O)-
12	4-bromophenyl	3-ethoxyphenyl	-CO-
13	4-ethylphenyl	2-methoxyphenyl	-C(O)-
14	4-styrenyl	3-tolyl	-C(O)-
15	4-isopropylphenyl	4-tolyl	-trans-CHCHC(O)-
16	4-isopropylphenyl	4-chlorophenyl	-trans-CHCHC(O)-
17	4-n-propylphenyl	phenyl	-trans-CHCHC(O)-
18	2-dimethylamino-6-naphthyl	4-ethylphenyl	-C(O)-
19	4-n-butylphenyl	3-methoxyphenyl	-C(O)-
20	4-t-butylphenyl	4-tolyl	-C(O)-
21	4-n-pentylphenyl	3-methoxyphenyl	-C(O)-
22	4-ethylphenyl	phenyl	-cyclopropyl-C(O)-
23	4-ethylphenyl		-C(O)-
24	4-ethylphenyl		-C(O)-



25	<i>trans</i> -2-(4-chlorophenyl)ethenyl	3-methoxyphenyl	-C(O)-
26	4-chlorophenylethynyl	3-methoxyphenyl	-C(O)-
27	phenylethynyl	3-methoxyphenyl	-C(O)-
28	4-ethylphenyl	phenyl	-C≡CC(O)-
29	4-ethylphenyl	phenyl	-CH <sub>2</sub> CH <sub>2</sub> C(O)-
30	3-tolyl	phenyl	-CH <sub>2</sub> CH <sub>2</sub> C(O)-
31	4-cyanophenyl	phenyl	-CH <sub>2</sub> CH <sub>2</sub> C(O)-
32	4- <i>t</i> -butylphenyl	3-methoxyphenyl	-C(O)-
33	4-ethylphenyl	3-chlorophenyl	-C(O)-
34	4-ethylphenyl	phenyl	-NHC(O)-
35	4-ethylphenyl	phenyl	- <i>trans</i> -CHCHC(O)-
36	4-ethylphenyl	phenylethynyl	-C(O)-
37	4-ethylphenyl	4-methoxyphenyl	-C≡CC(O)-
38	4- <i>n</i> -propylphenyl	2-methoxyphenyl	-C(O)-
39	4-ethylphenyl	4-ethylphenyl	-C(O)-
40	4-isopropylphenyl	4-ethylphenyl	-C(O)-
41	4- <i>n</i> -propylphenyl	4-ethylphenyl	-C(O)-
42	4-ethylphenyl	3-tolyl	-C(O)-
43	4-biphenyl	3,5-dimethoxyphenyl	-C(O)-
44	4- <i>n</i> -propylphenyl	3-tolyl	-C(O)-
45	4-ethylphenyl	3-ethoxyphenyl	-C(O)-
46	4-isopropylphenyl	3-ethoxyphenyl	-C(O)-
47	4- <i>n</i> -propylphenyl	3-ethoxyphenyl	-C(O)-
48	2-naphthyl	4-methoxyphenyl	-C(O)-
49	4-isopropylphenyl	phenyl	- <i>trans</i> -CHCHC(O)-
50	4-methoxyphenyl	phenyl	- <i>trans</i> -CHCHC(O)-
51	3-tolyl	phenyl	- <i>trans</i> -CHCHC(O)-
52	phenyl	phenyl	- <i>trans</i> -CHCHC(O)-
53	<i>trans</i> -2-(4-chlorophenyl)ethenyl	phenyl	- <i>trans</i> -CHCHC(O)-
54	<i>trans</i> -2-(4-chlorophenyl)ethenyl	4-tolyl	-C≡CC(O)-
55	3-tolyl	2,2-diphenethyl	-C(O)-
56	phenyl	2,2-diphenethyl	-C(O)-
57	4-methoxy-2,6-dimethylphenyl	3-methoxyphenyl	-C(O)-
58	4-ethylphenyl	phenyl-NH-	-CH <sub>2</sub> C(O)-
59	5-chloro-2-naphthyl	4-methoxyphenyl	-C(O)-
60	4-ethylphenyl	4-trifluoromethoxy phenyl	-CH <sub>2</sub> -
61	3,4-dichlorophenyl	3-trifluoromethoxy phenyl	-C(O)-




62	3-tolyl	3-trifluoromethoxy phenyl	-C(O)-
63	<i>trans</i> - $\beta$ -styrenyl	3-trifluoromethoxy phenyl	-C(O)-
64	4-bromophenyl	3-methoxyphenyl	-C(O)-
65	4-nitrophenyl	3-methoxyphenyl	-C(O)-
66	4-chlorophenylethynyl	4-tolyl	-C $\equiv$ CC(O)-
67	4-chlorophenylethynyl	phenyl	-C $\equiv$ CC(O)-
68	4-chlorophenylethynyl	4-tolyl	- <i>trans</i> -CHCHC(O)-
69	4-chlorophenylethynyl	phenyl	- <i>trans</i> -CHCHC(O)-
70	benzyl	3-methoxyphenyl	-C(O)-
71	benzyl	4-ethylphenyl	-C(O)-
72	4-nitrophenyl	2-methoxyphenyl	-C(O)-
73	4-n-propylphenyl	4-methoxyphenyl	-C(O)-
74	4-ethylphenyl	3-acetylphenyl	-C(O)-
75	4-acetylphenyl	4-tolyl	-C(O)-
76	4-tolyl	3-methoxyphenyl	-C(O)-
77	4-tolyl	4-ethylphenyl	-C(O)-
78	4-methoxyphenyl	3-tolyl	-C(O)-
79	phenyl	3-methoxyphenyl	-C(O)-
80	4-methoxyphenyl	3-methoxyphenyl	-C(O)-
81	2-thienyl	3-methoxyphenyl	-C(O)-
82	4-ethylphenyl	phenyl	-CH <sub>2</sub> C(O)-
83	4-n-butylphenyl	phenyl	-CH <sub>2</sub> C(O)-
84	4-n-pentylphenyl	phenyl	-CH <sub>2</sub> C(O)-
85	4-n-propylphenyl	4-nitrophenyl	-C(O)-
86	phenyl	4-ethylphenyl	-C(O)-
87	phenyl	3,4-dimethylphenyl	-C(O)-
88	3-trifluoromethylphenyl	3,4-dimethylphenyl	-C(O)-
89	4-ethylphenyl	3-tolyl	-C(O)-
90	4-isopropylphenyl	3-tolyl	-C(O)-
91	4-methoxyphenyl	3-tolyl	-C(O)-
92	4-ethylphenyl	4-fluorophenyl	-C(O)-
93	4-isopropylphenyl	4-fluorophenyl	-C(O)-
94	4-n-propylphenyl	4-fluorophenyl	-C(O)-
95	3-tolyl	4-ethylphenyl	-C(O)-
96	3-nitrophenyl	3,5-dimethoxyphenyl	-C(O)-
97	3-tolyl	3-tolyl	-C(O)-
98	4-ethylphenyl	diphenethyl	-C(O)-













99	4-isopropylphenyl	diphenethyl	-C(O)-
100	4-n-propylphenyl	diphenethyl	-C(O)-
101	4-methoxyphenyl	diphenethyl	-C(O)-
102	3-tolyl	phenoxy	-CH <sub>2</sub> C(O)-
103	4-ethylphenyl	2-thienyl	-C(O)-
104	4-ethylphenyl	4-dimethyl aminophenyl	-C(O)-
105	4-isopropylphenyl	2,2-dimethylethenyl	-C(O)-
106	4-isopropylphenyl	4-nitrophenyl	-C(O)-
107	3-tolyl	4-fluorophenyl	-C(O)-
108	3-tolyl	3-methoxyphenyl	-C(O)-
109	2,3,6-trimethyl-4-methoxyphenyl	3-methoxyphenyl	-C(O)-
110	4-methoxy-2,3,6-trimethylphenyl	phenyl	-C(O)-
111	2-phenethyl	3-methoxyphenyl	-C(O)-
112	<i>trans</i> -2-phenylethenyl	3-methoxyphenyl	-C(O)-
113	4-n-propylphenyl	3,5-dimethoxyphenyl	-C(O)-
114	4-t-amylphenyl	3-methoxyphenyl	-C(O)-
115	4-isopropylphenyl	4-methoxyphenyl	-CH <sub>2</sub> C(O)-
116	4-trifluoro methoxyphenyl	phenyl	-C(O)-
117	4-trifluormethylphenyl	3-tolyl	-C(O)-
118	3-chlorophenyl	3-methoxyphenyl	-C(O)-

Table 1B



Entry	R <sup>1</sup>	R <sup>2</sup>	R <sup>4</sup>	R <sup>3</sup>	-Y <sup>1</sup> -X <sup>1</sup> -
119	4-ethylphenyl	3-methoxyphenyl	benzyl	H	-C(O)-
120	4-ethylphenyl	cyclopropyl	benzyl	H	-C(O)-
121	4-ethylphenyl	3-methoxyphenyl	butyl	H	-C(O)-
122	4-ethylphenyl	3-methoxyphenyl		H	-C(O)-
123	4-ethylphenyl	cyclopropyl		H	-C(O)-
124	4-methoxy-2,3,6-trimethylphenyl	3-pyridine	H	H	-CH <sub>2</sub> -



					
125	4-ethylphenyl	cyclopropyl		H	-C(O)-
126	4-trifluoromethoxyphenyl	3-tolyl		H	-C(O)-
127	4-ethylphenyl	3-methoxyphenyl		H	-C(O)-
128	4-ethylphenyl	4-pyridine	H	H	-CH2-
129	4-methoxy-2,3,6-trimethylphenyl	2-pyridine	H	H	-CH2-
130	4-ethylphenyl	3-methoxyphenyl	ethyl	H	-C(O)-
131	4-ethylphenyl	tert-butyl	H		-C(O)-
132	4-ethylphenyl	3-methoxyphenyl	H		-C(O)-
133	4-ethylphenyl	3-methoxyphenyl	H	benzyl	-C(O)-
134	4-ethylphenyl	3-methoxyphenyl	H	butyl	-C(O)-
135	4-ethylphenyl		H	H	-C(O)-
136	4-ethylphenyl		H	H	-C(O)-
137	4-ethylphenyl		H	H	-C(O)-
138	4-ethylphenyl		H	H	-C(O)-
139	4-ethylphenyl	phenyl-NH-	H	H	-CH2C(O)-
140	4-ethylphenyl	benzyl-NH-	H	H	-CH2C(O)-
141	4-ethylphenyl	3,4-dimethoxybenzyl-NH-	H	H	-CH2C(O)-
142	4-ethylphenyl	3,4-dimethylphenyl-N(CH3)-	H	H	-CH2C(O)-
143	4-ethylphenyl	4-chlorobenzyl-N(CH3)-	H	H	-CH2C(O)-
144	4-ethylphenyl	benzyl-N(CH3)-	H	H	-CH2C(O)-
145	4-ethylphenyl	4-methoxyphenyl-N(CH3)-	H	H	-CH2C(O)-











					
146	4-ethylphenyl	3,4-dimethoxy phenyl-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	-CH <sub>2</sub> C(O)-
147	4-ethylphenyl		H	H	-CH <sub>2</sub> C(O)-
148	<i>para</i> -ethylphenyl		H	H	-CH <sub>2</sub> C(O)-
149	4-ethylphenyl		H	H	-C(O)-
150	4-ethylphenyl		H	H	-C(O)-
151	4-ethylphenyl		H	H	-C(O)-
152	4-ethylphenyl		H	H	-C(O)-
153	4-ethylphenyl	4-methoxyphenyl	ethyl	methyl	-C(O)-
154	4-ethylphenyl	4-methoxyphenyl		methyl	-C(O)-
155	4-ethylphenyl	3-pyridine	ethyl	H	-CH <sub>2</sub> -
156	4-ethylphenyl	3-methoxyphenyl	methyl	methyl	-CH <sub>2</sub> -
157	4-ethylphenyl	phenyl	methyl	H	-CH <sub>2</sub> CH <sub>2</sub> -

Table 1C

					
Entry	R <sup>1</sup>	R <sup>2</sup>	R <sup>4</sup>	R <sup>3</sup>	-Y <sup>1</sup> -X <sup>1</sup> -
158	4-ethylphenyl	3-methoxyphenyl	H	H	-C(O)-
159	4-ethylphenyl	4-methoxyphenyl	H	H	-C(O)-
160	4-ethylphenyl	t-butyl	H	H	-C(O)-



161	4-ethylphenyl	3-chlorophenyl	H	H	-C(O)-
162	4-ethylphenyl	cyclopropane		H	-C(O)-
163	4-ethylphenyl	methyl		H	-C(O)-
164	4-ethylphenyl	t-butyl		H	-C(O)-
165	4-ethylphenyl	3-methoxyphenyl		H	-C(O)-
166	4-ethylphenyl	3-methoxyphenyl	H		-C(O)-
167	4-ethylphenyl	cyclopropane	H		-C(O)-
168	4-ethylphenyl	t-butyl	H		-C(O)-
169	4-ethylphenyl	H		H	-CH2-
170	4-methoxy-2,3,6-trimethylphenyl	3-pyridyl	H	H	-CH2-
171	4-ethylphenyl	4-pyridyl	H	H	-CH2-
172	4-ethylphenyl	3-pyridyl	methyl	H	-CH2-
173	4-ethylphenyl	4-CF3O-phenyl	H	H	-CH2-
174	4-ethylphenyl	3-pyridyl	H		-CH2-
175	4-ethylphenyl	4-ethylphenyl	H		-CH2-

### EXAMPLES: BioAssays

#### <sup>86</sup>Rb Efflux Assays:

Cells stably transfected with cDNA for human Kv1.5 (in pcDNA3 vector) were grown as confluent monolayers in 96 well tissue culture plates in MEM alpha with 10% heat inactivated fetal bovine serum and 400 µg/ml G418. Cells were

incubated overnight in growth media containing 1  $\mu\text{Ci/ml}$   $^{86}\text{Rb}$  to permit intracellular uptake of the isotope. At the end of the incubation period, the  $^{86}\text{Rb}$  solution was aspirated and the cells washed three times with Earls Balanced Salt Solution (EBSS) which contained (in mM) 132 NaCl, 5.4 KCl, 1.8  $\text{CaCl}_2$ , 0.8 mM  $\text{MgCl}_2$ , 10 mM HEPES and 5 mM glucose. The cells were then preincubated for 10 minutes at room temperature in 100  $\mu\text{l}$ /well of EBSS or EBSS containing test compounds. At the end of this period the wells were aspirated and to each well was then added 100  $\mu\text{l}$  of a modified EBSS solution containing 70 mM KCl (NaCl replaced by KCl) and the compound to be tested. The high KCl concentration was utilized to depolarize the cells to membrane potentials that would activate Kv1.5 channels. After a 1 minute incubation in 70 mM KCl EBSS plus test compound, the solution was removed and placed into the appropriate well of a 96 well counting plate for analysis. Finally 100  $\mu\text{l}$  of 0.1% sodium dodecyl sulfate in EBSS was added to each well to lyse the cells. The lysate was taken for analysis to determine final cell content of  $^{86}\text{Rb}$ . Samples were counted in a Wallac Microbeta liquid scintillation counter by Cerenkov emission. Efflux was expressed as a percentage of the initial cell content of  $^{86}\text{Rb}$ .

The testing results of selective compounds from Tables 1A-C using this assay are reported in Table 2 (flux) as the potency for inhibition of  $^{86}\text{Rb}$  efflux through Kv1.5 potassium channels expressed in CHO cells by compounds of the invention.

#### Electrophysiological studies

Electrophysiological recordings of potassium currents in Chinese hamster ovary cells stably expressing the gene construct for the Kv1.5 potassium channel subunit were performed using the whole cell configuration of the patch clamp technique (Hamill et al., *Pflugers Archiv* 391:85, 1981). Cell lines expressing Kv1.5 were prepared using standard techniques known to those skilled in the art. Cells were plated on glass coverslips at a density of  $2 \times 10^4$  cells/coverslip and used within 24-48 hours. Solutions used for electrophysiological recordings were as follows. Extracellular bathing solutions contained (in mM) 132 NaCl, 5.4 KCl, 1.8  $\text{CaCl}_2$ , 0.8  $\text{MgCl}_2$ , 10 HEPES, 5 glucose at pH 7.3. Electrode pipette solutions for measuring

Kv1.5 contain (in mM) 100 KF, 40 KCl, 5 NaCl, 2 MgCl<sub>2</sub>, 5 mM EGTA, 10 mM HEPES and 5 glucose at pH 7.4, 295 mOsm. The coverslips were placed in a small chamber (volume ~ 200  $\mu$ l) on the mechanical stage of an inverted microscope and perfused (2 ml/min) with extracellular recording solution. Drug was applied using a series of narrow-bore glass capillary tubes (inner diameter ~100  $\mu$ m) positioned approximately 200  $\mu$ m from the cell.

The testing results of selective compounds from Tables 1A-C using this assay are reported in Table 2 (EP) as the potency for inhibition of Kv1.5 potassium currents by compounds of the invention.

Table 2

Entry #	IC <sub>50</sub> ( $\mu$ M) (EP)	IC <sub>50</sub> ( $\mu$ M) (flux)
1	0.25	6.8
13	0.4	>50
19	0.05	2.9
24	0.6	5.9
28	0.09	5.9
40	ND	9
60	0.5	>50
70	2.1	29
85	ND	46
97	ND	39
103	ND	20
110	ND	12
123	0.1	ND
132	0.5	ND
135	0.1	ND
158	0.6	ND
162	0.2	ND

The principles, preferred embodiments and modes of operation of the present invention have been described in the foregoing specification. The

invention which is intended to be protected herein, however, is not to be construed as limited to the particular forms disclosed, since they are to be regarded as illustrative rather than restrictive. Variations and changes may be made by those skilled in the art without departing from the spirit of the invention. Those skilled in the art will recognize variations in the processes as described above and will recognize appropriate modifications based on the above disclosure for making and using the compounds of the invention.

In the forgoing specification, the following abbreviations are used:


	<u>Designation</u>	<u>Reagent or Fragment</u>
10	m-CPBA	<i>meta</i> -chloroperoxybenzoic acid
	THF	tetrahydrofuran
	TLC	Thin Layer Chromotography
	DMF	dimethylformamide
	DMAP	<i>para</i> -dimethylaminopyridine
15	Me	methyl
	Et	ethyl
	EtOH	ethanol
	MeOH	methanol
	EtOAc	ethyl acetate
20	TsOH • H <sub>2</sub> O	<i>para</i> -toluenesulfonic acid • water
	NEt <sub>3</sub>	triethylamine
	DMSO	dimethylsulfoxide
	<i>n</i> -Pr	<i>n</i> -propyl
	NMR	nuclear magnetic resonance
25	MHz	megahertz
	Hz	hertz
	CDCl <sub>3</sub>	chloroform- <i>d</i>

<u>Designation</u>	<u>Reagent or Fragment</u>
UV	ultra-violet
$R_f$	retention factor
cat.	Catalytic

## CLAIMS

We claim:

1. A compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when R<sup>1</sup> is an optionally substituted aryl, then R<sup>1</sup> is not a dialkoxyphenyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),



alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,  
 5 CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 10 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 15 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an  
 20 optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;


25 X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted


amino); and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^4-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$ .

2. The compound of claim 1 wherein A and B are each H and the formula (I) has a stereochemical configuration of substituents in accordance with the following formula (Ia):



3. A compound of formula (II) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein  $t$  is 1, or 2;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted

heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when  $R^1$  is an optionally substituted aryl, then  $R^1$  is not a dialkoxyphenyl;

$Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ ,  $w$  is 0, 1, or 2 and  $q$  is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$  then  $X^2$  is not  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl;

$Z$  is H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2, or 3; and  $o$  is 0, 1, or 2


$X^1$  is  $C=O$ ,  $C=S$ ,  $SO_2$  or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl; and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is

not H, and (ii) that if  $R^2$  is  $R^a-O-$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$ .

4. A compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein  $t$  is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when  $R^1$  is an optionally substituted aryl, then  $R^1$  is not a dialkoxyphenyl;

$Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ ,  $w$  is 0, 1, or 2 and  $q$  is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$ , then  $X^2$  is not  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),

alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,  
 5 CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N- ;  
 10 where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are  
 15 attached form a heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;  
 20


X<sup>1</sup> is C=O, C=S, SO<sub>2</sub>, or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted  
 25

amino); and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^4-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$ .

5. The compound of claim 4 wherein A and B are each H and the formula (I) has a stereochemical configuration of substituents in accordance with the following formula (Ia):



6. The compound of claim 1 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

A and B are each H;

$Y^2$  is  $(CH_2)_q$ ,  $HC=CH$ , or ethynyl and  $q$  is 0, 1 or 2;

$X^2$  is  $SO_2$ ;

$R^1$  is selected from the group of an optionally substituted aryl and an optionally substituted heteroaryl;

$X^1$  is  $C=O$ ,  $C=S$ , or  $(CH_2)_n$ ; wherein  $n$  is 0, 1, or 2; and

$Z$  is H or  $OR^{14}$ .

7. The compound of claim 1 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

A and B are each H;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0, 1, or 2;

$X^2$  is  $SO_2$ ;

$R^1$  is H or an optionally substituted aryl selected from the group of phenyl and naphthyl;

5  $X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2; and

$Y^1$  is  $(CH_2)_p$ ,  $CH=CH$ , or ethynyl where  $p$  is 0, 1, 2 or 3.

8. The compound of claim 4 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

10  $t$  is 1;

$A$  and  $B$  are each H;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0;

$X^2$  is  $SO_2$ ;

15  $R^1$  is an optionally substituted aryl selected from the group of phenyl and naphthyl;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2; and

$Z$  is H, or OH; and

$Y^1$  is  $CH=CH$ , ethynyl or  $(CH_2)_p$ , where  $p$  is 0, 1, 2 or 3.

20 9. The compound of claim 8 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein  $R^3$  is H.

10. The compound of claim 9 wherein  $R^1$  is an optionally substituted phenyl.

25 11. A pharmaceutical composition comprising a compound of claim 1 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

12. A pharmaceutical composition comprising a compound of claim 2 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or

prodrug and a pharmaceutically acceptable diluent or carrier.

13. A pharmaceutical composition comprising a compound of claim 3 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

14. A pharmaceutical composition comprising a compound of claim 4 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

15. A pharmaceutical composition comprising a compound of claim 5 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof and a pharmaceutically acceptable diluent or carrier.

16. A pharmaceutical composition comprising a compound of claim 6 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

17. A pharmaceutical composition comprising a compound of claim 7 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.


18. A pharmaceutical composition comprising a compound of claim 8 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

19. A pharmaceutical composition comprising a compound of claim 9 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.



20. A pharmaceutical composition comprising a compound of claim 10 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

21. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted

amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl),  
 alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),  
 alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-  
 5 NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup>  
 or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or  
 C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,  
 CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a  
 counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or  
 10 alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally  
 substituted aryl, an optionally substituted aralkyl, an optionally substituted  
 heteroaryl, an optionally substituted heterocyclyl, an optionally substituted  
 heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-;  
 15 where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an  
 optionally substituted aryl, an optionally substituted aralkyl, an optionally  
 substituted heteroaryl, an optionally substituted heterocyclyl, an optionally  
 substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is  
 selected from the group consisting of H, alkyl, an optionally substituted aryl, an  
 20 optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally  
 substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally  
 substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are  
 attached form a heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is  
 25 selected from the group consisting of an optionally substituted aryl, an optionally  
 substituted heteroaryl, an optionally substituted heterocyclyl and an optionally  
 substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted

aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and


with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^4-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$ .

22. The method of claim 21 wherein the potassium channel is a voltage gated potassium channel.

23. The method of claim 22 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kur}$  potassium current.

24. The method of claim 22 wherein the potassium channel is Kv1.5.

25. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug:



wherein  $t$  is 1, or 2;

A and B are each H;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$ ,  $HC=CH$ , or ethynyl and  $q$  is 0, 1, or 2, with the proviso that when

$Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;

$X^1$  is  $C=O$ ,  $C=S$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and


with the provisos (i) that when  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, and (ii) that if  $R^2$  is  $R^A-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$ .

26. The method of claim 25 wherein the potassium channel is a voltage gated potassium channel.

27. The method of claim 26 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kw}$  potassium current.

28. The method of claim 26 wherein the potassium channel is Kv1.5.

29. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug:



wherein t is 1, or 2;

A and B are each H;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q is 0, 1, or 2, with the proviso that when q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an

optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;


$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and with the proviso that when  $Y^1$  is  $(CH_2)_p$  and  $p$  is 0, then  $R^2$  is not H.

30. The method of claim 29 wherein the potassium channel is a voltage gated potassium channel.

31. The method of claim 30 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kur}$  potassium current.

32. The method of claim 30 wherein the potassium channel is Kv1.5.

33. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1;

A and B are each H;

R<sup>1</sup> is alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q is 0;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an



optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the proviso that when  $Y^1$  is  $(CH_2)_p$  and  $p$  is 0, then  $R^2$  is not H.


34. The method of claim 33 wherein the potassium channel is a voltage gated potassium channel.

35. The method of claim 34 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kw}$  potassium current.

36. The method of claim 34 wherein the potassium channel is Kv1.5.

37. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate,

hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,

$\text{CH}_2\text{N}(\text{R}^9)_3\text{L}$ , or  $\text{CO}_2\text{R}^9$ ; each  $\text{R}^9$  is independently selected from H, or alkyl; L is a counter ion;  $\text{R}^{15}$  is H, or alkyl; and  $\text{R}^{16}$  is H, alkyl or  $\text{CO}_2\text{R}^{10}$  and  $\text{R}^{10}$  is H, or alkyl;

$\text{R}^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $\text{R}^a\text{-O-}$ , and  $\text{R}^b\text{R}^c\text{-N-}$ ; where  $\text{R}^a$  and  $\text{R}^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $\text{R}^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $\text{R}^b$  and  $\text{R}^c$  along with the nitrogen to which they are attached form a heterocyclyl;


$\text{Y}^1$  is  $(\text{CH}_2)_p$ ,  $\text{CHR}^{17}(\text{CH}_2)_o$ ,  $\text{HC}=\text{CH}$ , or ethynyl; where  $\text{R}^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

$\text{X}^1$  is  $\text{C}=\text{O}$ ,  $\text{C}=\text{S}$ ,  $\text{SO}_2$  or  $(\text{CH}_2)_n$ ; where n is 0, 1, or 2;

$\text{R}^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if  $\text{Y}^1$  is  $(\text{CH}_2)_p$ , p is 0 and  $\text{X}^1$  is not  $(\text{CH}_2)_n$ , then  $\text{R}^2$  is not H, (ii) that if  $\text{R}^2$  is  $\text{R}^a\text{-O}$  and  $\text{Y}^1$  is  $(\text{CH}_2)_p$  with p=0, then  $\text{X}^1$  is not  $\text{SO}_2$  and (iii) if Z is not H,  $\text{OR}^{14}$ ,  $\text{SR}^{14}$  or  $\text{NR}^{15}\text{R}^{16}$ , then  $\text{X}^2$  must be  $\text{SO}_2$ .

38. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formula (II) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, HC=CH, or ethynyl and q is 0, 1, or 2, with the proviso that when Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or

$C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;


$X^1$  is  $C=O$ ,  $C=S$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that when  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, and (ii) that if  $R^2$  is  $R^a-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not

SO<sub>2</sub>.

39. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formula (III) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q is 0, 1, or 2, with the proviso that when q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup>

or  $\text{NR}^{15}\text{R}^{16}$ ; where  $\text{R}^{14}$  is selected from the group consisting of H,  $(\text{CH}_2)_m\text{-R}^8$ , or  $\text{C(O)}\text{-(CH}_2)_r\text{-R}^8$ ; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3;  $\text{R}^8$  is  $\text{CH}_2\text{N(R}^9)_2$ ,  $\text{CH}_2\text{N(R}^9)_3\text{L}$ , or  $\text{CO}_2\text{R}^9$ ; each  $\text{R}^9$  is independently selected from H, or alkyl; L is a counter ion;  $\text{R}^{15}$  is H, or alkyl; and  $\text{R}^{16}$  is H, alkyl or  $\text{CO}_2\text{R}^{10}$  and  $\text{R}^{10}$  is H, or alkyl;

$\text{R}^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $\text{R}^a\text{-O-}$ , and  $\text{R}^b\text{R}^c\text{-N-}$ ; where  $\text{R}^a$  and  $\text{R}^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $\text{R}^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $\text{R}^b$  and  $\text{R}^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$\text{Y}^1$  is  $(\text{CH}_2)_p$ ,  $\text{CHR}^{17}(\text{CH}_2)_o$ ,  $\text{HC=CH}$ , or ethynyl; where  $\text{R}^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;


$\text{X}^1$  is  $\text{C=O}$ , or  $(\text{CH}_2)_n$ ; where n is 0, 1, or 2;

$\text{R}^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the proviso that when  $\text{Y}^1$  is  $(\text{CH}_2)_p$  and p is 0, then  $\text{R}^2$  is not H.

40. A method for treating cardiac arrhythmias which comprises administering to a

patient in need thereof, a pharmaceutically effective amount of a compound of formula (IV) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof.



wherein  $t$  is 1;

**A and B are each H:**

R<sup>1</sup> is alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0;

**X<sup>2</sup> is SO<sub>2</sub>;**

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a



counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;


$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and with the proviso that when  $Y^1$  is  $(CH_2)_p$  and  $p$  is 0, then  $R^2$  is not H.

## AMENDED CLAIMS

[received by the International Bureau on 7 July 1999 (07.07.99);  
original claims 1-40 replaced by new claims 1-41(31 pages)]

1. A compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when R<sup>1</sup> is an optionally substituted aryl, then R<sup>1</sup> is not a dialkoxyphenyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted

heteroaralkyl; an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),  
 5 alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a  
 10 counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted  
 15 heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N- ; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is  
 20 selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;

25 Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally


substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

$X^1$  is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;


$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if  $Y^1$  is (CH<sub>2</sub>)<sub>p</sub>, p is 0 and  $X^1$  is not (CH<sub>2</sub>)<sub>n</sub>, then  $R^2$  is not H, (ii) that if  $R^2$  is R<sup>a</sup>-O and  $Y^1$  is (CH<sub>2</sub>)<sub>p</sub> with p=0, then  $X^1$  is not SO<sub>2</sub>, (iii) if Z is not H, OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>, then  $X^2$  must be SO<sub>2</sub> and (iv) if t=1, A, B and Z are each H,  $X^2$  is C=O, or C=S,  $Y^2$  is NH,  $R^3$  is H or alkyl,  $R^1$  is H or alkyl,  $X^1$  is C=O, or C=S,  $R^4$  is H, and  $Y^1$  is (CH<sub>2</sub>)<sub>p</sub> with p=0 then  $R^2$  cannot be alkyl, or R<sup>b</sup>R<sup>c</sup>-N.

2. The compound of claim 1 wherein A and B are each H and the formula (I) has a stereochemical configuration of substituents in accordance with the following formula (Ia):



3. A compound of formula (II) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when  $R^1$  is an optionally substituted aryl, then  $R^1$  is not a dialkoxyphenyl;

$Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ , w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$  then  $X^2$  is not  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl;

Z is H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl; L is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an

optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;


$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2, or 3; and  $o$  is 0, 1, or 2

$X^1$  is  $C=O$ ,  $C=S$ ,  $SO_2$  or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, or an optionally substituted heteroaryl; and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^a-O-$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and (iii) if  $t=1$ ,  $Z$  is H,  $X^2$  is  $C=O$ , or  $C=S$ ,  $Y^2$  is NH,  $R^3$  is H or alkyl,  $R^1$  is H or alkyl,  $X^1$  is  $C=O$ , or  $C=S$ ,  $R^4$  is H, and  $Y^1$  is  $(CH_2)_p$  with  $p=0$  then  $R^2$  cannot be alkyl, or  $R^bR^c-N$ .

4. A compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

5  $R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when  $R^1$  is an optionally substituted aryl, then  $R^1$  is not a dialkoxyphenyl;

10  $Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ , w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$ , then  $X^2$  is not  $SO_2$ ;

15  $R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

20 Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl; L is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or  
25 alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted

heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;


$X^1$  is  $C=O$ ,  $C=S$ ,  $SO_2$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, (ii) that if  $R^2$  is  $R^a-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$ , (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$  and (iv) if  $t=1$ , A, B and Z are each H,  $X^2$  is  $C=O$ , or  $C=S$ ,  $Y^2$  is NH,  $R^3$  is H or alkyl,  $R^1$  is H or alkyl,  $X^1$  is  $C=O$ , or  $C=S$ ,  $R^4$  is H, and  $Y^1$  is  $(CH_2)_p$  with  $p=0$  then  $R^2$  cannot be alkyl, or  $R^bR^c-N$ .



5. The compound of claim 4 wherein A and B are each H and the formula (I) has a stereochemical configuration of substituents in accordance with the following formula (Ia):



6. The compound of claim 1 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

A and B are each H;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, HC=CH, or ethynyl and q is 0, 1 or 2;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>1</sup> is selected from the group of an optionally substituted aryl and an optionally substituted heteroaryl;

X<sup>1</sup> is C=O, C=S, or (CH<sub>2</sub>)<sub>n</sub>; wherein n is 0, 1, or 2; and Z is H or OR<sup>14</sup>.

7. The compound of claim 1 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

A and B are each H;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q is 0, 1, or 2;

X<sup>2</sup> is SO<sub>2</sub>;

R<sup>1</sup> is H or an optionally substituted aryl selected from the group of phenyl

and naphthyl;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2; and

$Y^1$  is  $(CH_2)_p$ ,  $CH=CH$ , or ethynyl where  $p$  is 0, 1, 2 or 3.

5 8. The compound of claim 4 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein

$t$  is 1;

A and B are each H;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0;

10  $X^2$  is  $SO_2$ ;

$R^1$  is an optionally substituted aryl selected from the group of phenyl and naphthyl;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2; and

Z is H, or OH; and

15  $Y^1$  is  $CH=CH$ , ethynyl or  $(CH_2)_p$ , where  $p$  is 0, 1, 2 or 3.

9. The compound of claim 8 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof, wherein  $R^3$  is H.

10. The compound of claim 9 wherein  $R^1$  is an optionally substituted phenyl.

20 11. A pharmaceutical composition comprising a compound of claim 1 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

25 12. A pharmaceutical composition comprising a compound of claim 2 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

13. A pharmaceutical composition comprising a compound of claim 3 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

5 14. A pharmaceutical composition comprising a compound of claim 4 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

10 15. A pharmaceutical composition comprising a compound of claim 5 or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof and a pharmaceutically acceptable diluent or carrier.

15 16. A pharmaceutical composition comprising a compound of claim 6 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

20 17. A pharmaceutical composition comprising a compound of claim 7 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.


18. A pharmaceutical composition comprising a compound of claim 8 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

25 19. A pharmaceutical composition comprising a compound of claim 9 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or

prodrug and a pharmaceutically acceptable diluent or carrier.

20. A pharmaceutical composition comprising a compound of claim 10 or its pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug and a pharmaceutically acceptable diluent or carrier.

21. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

R<sup>1</sup> is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub>, (CH<sub>2</sub>)<sub>w</sub>O, HC=CH, ethynyl or NH, w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>q</sub> and q=0, then R<sup>1</sup> cannot be H;

X<sup>2</sup> is C=O, C=S, or SO<sub>2</sub>; with the proviso that if Y<sup>2</sup> is (CH<sub>2</sub>)<sub>w</sub>O, then X<sup>2</sup> is not SO<sub>2</sub>;

R<sup>3</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted

aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

5           Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or  
10           C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

15           R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N- ; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is  
20           selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are  
25           attached form a heterocyclyl;

          Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is

selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

$X^1$  is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

5  $R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and


10 with the provisos (i) that if Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, p is 0 and X<sup>1</sup> is not (CH<sub>2</sub>)<sub>n</sub>, then R<sup>2</sup> is not H, (ii) that if R<sup>2</sup> is R<sup>4</sup>-O and Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub> with p=0, then X<sup>1</sup> is not SO<sub>2</sub> and (iii) if Z is not H, OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>, then X<sup>2</sup> must be SO<sub>2</sub>.

22. The method of claim 21 wherein the potassium channel is a voltage gated potassium channel.

15 23. The method of claim 22 wherein the potassium channel is selected from a potassium channel responsible for cardiac I<sub>Kur</sub> potassium current.

24. The method of claim 22 wherein the potassium channel is Kv1.5.

25. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing  
20 said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug:



wherein t is 1, or 2;

A and B are each H;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$ ,  $HC=CH$ , or ethynyl and q is 0, 1, or 2, with the proviso that when  $Y^2$  is  $(CH_2)_q$  and q=0, then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl; L is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or

alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;

$X^1$  is  $C=O$ ,  $C=S$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that when  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, and (ii) that if  $R^2$  is  $R^a-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$ .




26. The method of claim 25 wherein the potassium channel is a voltage gated potassium channel.

27. The method of claim 26 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kur}$  potassium current.

28. The method of claim 26 wherein the potassium channel is Kv1.5.

29. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug:



wherein t is 1, or 2;

A and B are each H;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$  and q is 0, 1, or 2, with the proviso that when q=0, then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted

heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl),  
 alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),  
 5 alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-  
 NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup>  
 or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or  
 C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,  
 CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a  
 10 counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or  
 alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted  
 aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an  
 15 optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are  
 independently selected from the group consisting of alkyl, an optionally substituted  
 aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an  
 optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of  
 20 H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an  
 optionally substituted heteroaryl, an optionally substituted heterocyclyl, a  
 optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  
 or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a  
 heterocyclyl;

25 Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is  
 selected from the group consisting of an optionally substituted aryl, an optionally  
 substituted heteroaryl, an optionally substituted heterocyclyl and an optionally

substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where n is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and


with the proviso that when  $Y^1$  is  $(CH_2)_p$  and p is 0, then  $R^2$  is not H.

30. The method of claim 29 wherein the potassium channel is a voltage gated potassium channel.

31. The method of claim 30 wherein the potassium channel is selected from a potassium channel responsible for cardiac  $I_{Kur}$  potassium current.

32. The method of claim 30 wherein the potassium channel is Kv1.5.

33. A method for inhibiting potassium transport across cellular membranes possessing potassium channels comprising exposing a cell membrane possessing said channels to the presence of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1;

A and B are each H;

$R^1$  is alkyl, or is selected from the group consisting of an optionally

substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an

optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;

5           Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

          X<sup>1</sup> is C=O, or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

10           R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

15           with the proviso that when Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub> and p is 0, then R<sup>2</sup> is not H.


34. The method of claim 33 wherein the potassium channel is a voltage gated potassium channel.

35. The method of claim 34 wherein the potassium channel is selected from a potassium channel responsible for cardiac I<sub>Kur</sub> potassium current.

20 36. The method of claim 34 wherein the potassium channel is Kv1.5.

37. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:

25



wherein t is 1, or 2;

A and B are each H, or taken together form a bond between the substituted carbons;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$ ,  $(CH_2)_wO$ ,  $HC=CH$ , ethynyl or  $NH$ , w is 0, 1, or 2 and q is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $C=O$ ,  $C=S$ , or  $SO_2$ ; with the proviso that if  $Y^2$  is  $(CH_2)_wO$ , then  $X^2$  is not  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,

$\text{CH}_2\text{N}(\text{R}^9)_3\text{L}$ , or  $\text{CO}_2\text{R}^9$ ; each  $\text{R}^9$  is independently selected from H, or alkyl; L is a counter ion;  $\text{R}^{15}$  is H, or alkyl; and  $\text{R}^{16}$  is H, alkyl or  $\text{CO}_2\text{R}^{10}$  and  $\text{R}^{10}$  is H, or alkyl;

$\text{R}^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $\text{R}^a\text{-O-}$ , and  $\text{R}^b\text{R}^c\text{-N-}$ ; where  $\text{R}^a$  and  $\text{R}^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $\text{R}^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $\text{R}^b$  and  $\text{R}^c$  along with the nitrogen to which they are attached form a heterocyclyl;

$\text{Y}^1$  is  $(\text{CH}_2)_p$ ,  $\text{CHR}^{17}(\text{CH}_2)_o$ ,  $\text{HC=CH}$ , or ethynyl; where  $\text{R}^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;


$\text{X}^1$  is  $\text{C=O}$ ,  $\text{C=S}$ ,  $\text{SO}_2$  or  $(\text{CH}_2)_n$ ; where n is 0, 1, or 2;

$\text{R}^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if  $\text{Y}^1$  is  $(\text{CH}_2)_p$ , p is 0 and  $\text{X}^1$  is not  $(\text{CH}_2)_n$ , then  $\text{R}^2$

is not H, (ii) that if  $R^2$  is  $R^A-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$  and  
 (iii) if  $Z$  is not H,  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ , then  $X^2$  must be  $SO_2$ .

38. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of  
 5 formual (II) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein  $t$  is 1, or 2;

10 A and B are each H;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

15  $Y^2$  is  $(CH_2)_q$ ,  $HC=CH$ , or ethynyl and  $q$  is 0, 1, or 2, with the proviso that when  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

20  $R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl),



alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl),  
 alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-  
 NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup>  
 or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or  
 5 C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>,  
 CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a  
 counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or  
 alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted  
 10 aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an  
 optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are  
 independently selected from the group consisting of alkyl, an optionally substituted  
 aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 15 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an  
 optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of  
 H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an  
 optionally substituted heteroaryl, an optionally substituted heterocyclyl, an  
 optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  
 20 or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a  
 heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is  
 selected from the group consisting of an optionally substituted aryl, an optionally  
 substituted heteroaryl, an optionally substituted heterocyclyl and an optionally  
 25 substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;


X<sup>1</sup> is C=O, C=S, or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted

aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that when  $Y^1$  is  $(CH_2)_p$ ,  $p$  is 0 and  $X^1$  is not  $(CH_2)_n$ , then  $R^2$  is not H, and (ii) that if  $R^2$  is  $R^4-O$  and  $Y^1$  is  $(CH_2)_p$  with  $p=0$ , then  $X^1$  is not  $SO_2$ .

39. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formual (III) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein  $t$  is 1, or 2;

A and B are each H;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$  and  $q$  is 0, 1, or 2, with the proviso that when  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted

aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

5 Z is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl), OR<sup>14</sup>, SR<sup>14</sup> or NR<sup>15</sup>R<sup>16</sup>; where R<sup>14</sup> is selected from the group consisting of H, (CH<sub>2</sub>)<sub>m</sub>-R<sup>8</sup>, or  
 10 C(O)-(CH<sub>2</sub>)<sub>r</sub>-R<sup>8</sup>; m is 1, 2, 3, or 4; r is 0, 1, 2, or 3; R<sup>8</sup> is CH<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, CH<sub>2</sub>N(R<sup>9</sup>)<sub>3</sub>L, or CO<sub>2</sub>R<sup>9</sup>; each R<sup>9</sup> is independently selected from H, or alkyl; L is a counter ion; R<sup>15</sup> is H, or alkyl; and R<sup>16</sup> is H, alkyl or CO<sub>2</sub>R<sup>10</sup> and R<sup>10</sup> is H, or alkyl;

R<sup>2</sup> is selected from the group consisting of H, alkyl, an optionally substituted  
 15 aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl, R<sup>a</sup>-O-, and R<sup>b</sup>R<sup>c</sup>-N-; where R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an  
 20 optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; R<sup>c</sup> is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  
 25 or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is


selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where n is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the proviso that when  $Y^1$  is  $(CH_2)_p$  and p is 0, then  $R^2$  is not H.

40. A method for treating cardiac arrhythmias which comprises administering to a patient in need thereof, a pharmaceutically effective amount of a compound of formula (IV) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



wherein t is 1;

A and B are each H;

$R^1$  is alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;

$Y^2$  is  $(CH_2)_q$  and q is 0;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or, alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or  $R^b$  and  $R^c$  along with the nitrogen to which they are attached form a

heterocyclyl;


$Y^1$  is  $(CH_2)_p$ ,  $CHR^{17}(CH_2)_o$ ,  $HC=CH$ , or ethynyl; where  $R^{17}$  is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl;  $p$  is 0, 1, 2 or 3; and  $o$  is 0, 1 or 2;

$X^1$  is  $C=O$ , or  $(CH_2)_n$ ; where  $n$  is 0, 1, or 2;

$R^4$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the proviso that when  $Y^1$  is  $(CH_2)_p$  and  $p$  is 0, then  $R^2$  is not H.

41. A compound of formula (I) or a pharmaceutically acceptable salt, ester, amide, complex, chelate, hydrate, stereoisomer, crystalline or amorphous form, metabolite, metabolic precursor or prodrug thereof:



A and B are each H, or taken together form a bond between the substituted carbons;

$R^1$  is H, alkyl, or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted

heterocyclyl and an optionally substituted carbocycloalkyl with the proviso that when  $R^1$  is an optionally substituted aryl, then  $R^1$  is not a dialkoxyphenyl;

$Y^2$  is  $(CH_2)_q$ ,  $HC=CH$ , ethynyl or  $NH$ ,  $w$  is 0, 1, or 2 and  $q$  is 0, 1, or 2, with the proviso that if  $Y^2$  is  $(CH_2)_q$  and  $q=0$ , then  $R^1$  cannot be H;

$X^2$  is  $SO_2$ ;

$R^3$  is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino);

$Z$  is H, alkyl, alkyenyl, alkylene(heterocyclyl), alkylene(heteroaryl), alkylene-NHC(O)(alkyl), alkylene-NHC(O)(aryl), alkylene-NHC(O)(heterocyclyl), alkylene-NHC(O)(heteroaryl), alkylene-NHC(O)-(alkylene-heterocyclyl), alkylene-NHC(O)-(heteroaralkyl), alkylene-C(O)(alkyl), alkylene-C(O)O(alkyl),  $OR^{14}$ ,  $SR^{14}$  or  $NR^{15}R^{16}$ ; where  $R^{14}$  is selected from the group consisting of H,  $(CH_2)_m-R^8$ , or  $C(O)-(CH_2)_r-R^8$ ;  $m$  is 1, 2, 3, or 4;  $r$  is 0, 1, 2, or 3;  $R^8$  is  $CH_2N(R^9)_2$ ,  $CH_2N(R^9)_3L$ , or  $CO_2R^9$ ; each  $R^9$  is independently selected from H, or alkyl;  $L$  is a counter ion;  $R^{15}$  is H, or alkyl; and  $R^{16}$  is H, alkyl or  $CO_2R^{10}$  and  $R^{10}$  is H, or alkyl;

$R^2$  is selected from the group consisting of H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, an optionally substituted carbocycloalkyl,  $R^a-O-$ , and  $R^bR^c-N-$ ; where  $R^a$  and  $R^b$  are independently selected from the group consisting of alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl;  $R^c$  is selected from the group consisting of

H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl, an optionally substituted heteroaralkyl, and an optionally substituted carbocycloalkyl; or R<sup>b</sup> and R<sup>c</sup> along with the nitrogen to which they are attached form a heterocyclyl;

Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, CHR<sup>17</sup>(CH<sub>2</sub>)<sub>o</sub>, HC=CH, or ethynyl; where R<sup>17</sup> is alkyl or is selected from the group consisting of an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted heterocyclyl and an optionally substituted carbocycloalkyl; p is 0, 1, 2 or 3; and o is 0, 1 or 2;

X<sup>1</sup> is C=O, C=S, SO<sub>2</sub> or (CH<sub>2</sub>)<sub>n</sub>; where n is 0, 1, or 2;

R<sup>4</sup> is H, alkyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteroaryl, an optionally substituted heteroaralkyl; an optionally substituted heterocycle, an optionally substituted heterocyclyl, an optionally substituted carbocycloalkyl, or an alkylene-(substituted amino); and

with the provisos (i) that if Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub>, p is 0 and X<sup>1</sup> is not (CH<sub>2</sub>)<sub>n</sub>, then R<sup>2</sup> is not H; (ii) that if R<sup>2</sup> is R<sup>a</sup>-O and Y<sup>1</sup> is (CH<sub>2</sub>)<sub>p</sub> with p=0, then X<sup>1</sup> is not SO<sub>2</sub>.



# INTERNATIONAL SEARCH REPORT

Inter      national Application No  
PCT/US 99/01663

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6    C07C311/20    C07C311/13    C07C311/28    C07D207/16    C07D213/40  
          C07D213/42    C07D217/26    C07D333/38    A61K31/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6    C07C    C07D    A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 1 479 544 A (AMERICAN CYANAMID) 13 July 1977 see page 1 - page 2; examples 2,5,7-9,58 ---	1,3,4
A	US 5 631 275 A (H. ENGLERT, ET AL.) 20 May 1997 see column 1 - column 7 ---	1-40
P, A	WO 98 04521 A (ICAGEN, ET AL.) 5 February 1998 see page 4 - page 5 -----	1-40

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 April 1999

Date of mailing of the international search report

07/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

English, R

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/01663

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 21-40  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 21-40  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

information on patent family members

Inter nal Application No

PCT/US 99/01663

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1479544 A	13-07-1977	US 3953506 A	27-04-1976
		US 3953606 A	27-04-1976
		DE 2505301 A	14-08-1975
		FR 2260339 A	05-09-1975
		JP 50108250 A	26-08-1975
		LU 71799 A	24-06-1975
		NL 7501474 A	11-08-1975
		US 4005140 A	25-01-1977
		BE 825275 A	06-08-1975
		ZA 7500416 A	28-01-1976
US 5631275 A	20-05-1997	DE 4344957 A	06-07-1995
		US 5731341 A	24-03-1998
		AU 679136 B	19-06-1997
		AU 8178994 A	06-07-1995
		CA 2138930 A	01-07-1995
		CN 1109872 A	11-10-1995
		EP 0661264 A	05-07-1995
		FI 946136 A	01-07-1995
		HU 70830 A	28-11-1995
		JP 7206807 A	08-08-1995
		NO 945083 A	03-07-1995
		NZ 270265 A	26-10-1995
		ZA 9410380 A	29-08-1995
WO 9804521 A	05-02-1998	AU 3803597 A	20-02-1998

**THIS PAGE BLANK (USPTO)**